

Teramycin Vet

Oxytetracycline Hydrochloride

Composition:

Teramycin Vet 500 Tablet: Each tablet contains Oxytetracycline Hydrochloride USP 500 mg.

Indication:

Recommended for the control and treatment of the following diseases of beef and dairy calves caused by organisms sensitive to oxytetracycline: bacterial enteritis (scours, diarrhea) caused by *Salmonella typhimurium* and *Escherichia coli* (colibacillosis); bacterial pneumonia (shipping fever complex, pasteurellosis) caused by *Pasteurella multocida*.

Dosage:

For the control of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally half a bolus (250 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (5 mg/lb body weight in divided doses) for up to four consecutive days.

For the treatment of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally one bolus (500 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (10 mg/lb body weight in divided doses) for up to four consecutive days.

Dosage should continue until the animal returns to normal and for 24 to 48 hours after symptoms have subsided. Treatment should not exceed four consecutive days.

Contraindication:

Not for use in lactating dairy cattle. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Precautions:

Exceeding the recommended dosage level of one bolus (500 mg Oxytetracycline hydrochloride) per 100 lb body weight every 12 hours, or administration at the recommended level for more than four consecutive days, may result in antibiotic residues beyond the withdrawal time.

Organisms may vary in their degree of susceptibility to any chemotherapy. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reexamined.

Rarely do side reactions or allergic manifestations occur in calves treated with Teramycin. If any unusual reactions are noted, discontinue use of the drug immediately and call a veterinarian.

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving Teramycin conjunction with penicillin.

Storage:

Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F).

Packaging:

Teramycin Vet 500 Tablet: Each pack contains 5x4's tablet.

Manufactured by



Ziska Pharmaceuticals Ltd.

Kaliakoir, Gazipur, Bangladesh